
Index

A

Akaike information criterion, 409–412

corrected, 412

generalized, 411–413

all-pole signals, 94

amplitude and phase estimation (APES)

method, 256–259, 303

for gapped data, 259–262

for spatial spectra, 319–326

for two-dimensional signals, 267–269

amplitude spectrum, 253, 258

Capon estimates of, 254–255

angle of arrival, 276

aperture, 276

APES method, *see* amplitude and phase
estimation method

apodization, 62–67

AR process, *see* autoregressive process

AR spectral estimation, *see* autoregressive
spectral estimation

ARMA process, *see* autoregressive moving
average process

array

aperture of, 276

beamforming resolution, 334, 335

beamspace processing, 337, 338

beamwidth, 290, 336

broadband signals in, 281

coherent signals in, 293, 340

isotropic, 336

L-shaped, 335

narrowband, 281, 283

planar, 276

uniform linear, 283–285

array model, 277–285

autocorrelation function, 122

autocorrelation method, 97

autocovariance sequence

computation using FFT, 58–60

computer generation of, 20

definition of, 5

estimates, 24

estimation variance, 76

extensions, 123–125, 182

for signals with unknown means, 75

for sinusoidal signals, 151, 152

generation from ARMA parameters, 136

mean-square convergence of, 179

of ARMA processes, 93–94

properties, 5–6

autoregressive (AR) process
 covariance structure, 93
 definition of, 92
 stability of, 139
 autoregressive moving average (ARMA)
 process
 covariance structure, 93
 definition of, 92
 multivariate, 113–122
 state–space equations, 114
 autoregressive moving-average spectral
 estimation, 107–122
 least-squares method, 111–113
 modified Yule–Walker method, 108–111
 multivariate, 117–122
 autoregressive spectral estimation, 94–99
 autocorrelation method, 97
 Burg method, 125–127
 covariance method, 94, 97
 least-squares method, 96–99
 postwindow method, 98
 prewindow method, 98
 Yule–Walker method, 94

B

backward prediction, 122, 123, 137
 bandpass signal, 277
 bandwidth
 approximate formula, 81
 definition of, 71
 equivalent, 43, 58, 73, 234
 Bartlett method, 52–53
 Bartlett window, 31, 45
 baseband signal, 278
 basis
 linearly parameterized, 203–209
 null space, 203–209
 Bayesian information criterion, 413–417
 beamforming, 288–291, 300–302
 beamforming method, 307
 and CLEAN, 326–331
 beamspace processing, 337, 338

beamwidth, 290, 335, 336
 BIC rule, 413–417
 Blackman window, 45
 Blackman–Tukey method, 39–42
 computation using FFT, 60–62
 nonnegativeness property, 41
 block–Hankel matrix, 118
 broadband signal, 281
 Burg method, 125–127

C

CAPES method, 259
 Capon method, 232–242, 302–306
 as a matched filter, 270
 comparison with APES, 258–259
 constrained, 311–319
 derivation of, 232–238, 270
 for damped sinusoids, 253–256
 for DOA estimation, 291–293
 for two-dimensional signals, 267–269
 relationship to AR methods, 238–242,
 246–249
 robust, 306–319
 spectrum of, 271
 stochastic signal, 302–303
 Carathéodory parameterization, 311
 carrier frequency, 277
 Cauchy–Schwartz inequality, 270, 291, 314,
 317, 330, 360–361
 for functions, 361
 for vectors, 360–361
 centrosymmetric matrix, 178, 332
 Chebyshev inequality, 211
 Chebyshev window, 44
 chi-squared distribution, 184
 Cholesky factor, 134, 358
 circular Gaussian distribution, 80, 331, 380,
 387
 circular white noise, 34, 39
 CLEAN algorithm, 326–331
 coherence spectrum, 67–70
 coherent signals, 293, 340

column space, 342
complex demodulation, 280
complex envelope, 280
complex modulation, 279
complex white noise, 34
concave function, 192
condition number, 212
 and AR parameter estimation, 109
 and forward–backward approach, 212
 definition of, 366
confidence interval, 79
consistent estimator, 373
consistent linear equations, 364–366
constant-modulus signal, 300–302
constrained Capon method, 311–319
continuous spectra, 90
convergence
 in probability, 211
 mean-square, 179, 180, 211
 uniform, 272
corrected Akaike information criterion, 412
correlation coefficient, 15
correlogram method, 24–27
covariance
 definition of, 5
 matrix, 6
covariance fitting, 303–306, 329
 using CLEAN, 328–331
covariance-fitting criterion, 132
covariance function, *see* autocovariance sequence
covariance matrix
 diagonalization of, 139
 eigenvalue decomposition of, 309, 315
 persymmetric, 178, 332
 properties of, 6
covariance method, 97
covariance sequence, *see* autocovariance sequence
covariance-fitting criterion, 132

Cramér–Rao bound, 373–396, 400
 for Gaussian distributions, 378–383
 for general distributions, 376–378
 for line spectra, 383–384
 for rational spectra, 384–386
 for spatial spectra, 387–396
 for unknown model order, 376
cross covariance sequence, 19
cross-spectrum, 13, 19, 67
cyclic minimization, 189–190
cyclic minimizer, 260

D
damped sinusoidal signals, 203–209, 253–256
Daniell method, 55–58
delay operator, 11
Delsarte–Genin Algorithm, 102–106
demodulation, 280
diagonal loading, 312–319
Dirac impulse, 152
direction of arrival, 276
direction-of-arrival estimation, 275–298
 beamforming, 288–291
 Capon method, 291–293
 ESPRIT method, 297–298
 Min–Norm method, 297
 MUSIC method, 297
 nonlinear least-squares method, 294
 nonparametric methods, 285–293
 parametric methods, 293–298
 Pisarenko method, 297
 Yule–Walker method, 295
direction vector uncertainty, 306–319
Dirichlet kernel, 33
discrete Fourier transform (DFT), 27
 linear transformation interpretation, 77
discrete signals, 2
discrete spectrum, 153
discrete-time Fourier transform (DTFT), 3

discrete-time system, 11
 finite-impulse response (FIR), 18
 frequency response, 221
 minimum phase, 93, 135
 transfer function, 221
 displacement operator, 129
 displacement rank, 130
 Doppler frequency, 334
 Durbin's method, 106, 113

E

efficiency, statistical, 375
 eigenvalue, 345
 of a matrix product, 347
 eigenvalue decomposition, 309, 315, 344–351
 eigenvector, 345
 EM algorithm, 188–194
 energy-spectral density, 3
 Capon estimates of, 254–255
 of damped sinusoids, 253
 ergodic, 179
 ESPRIT method
 and min–norm, 212, 213
 combined with HOYW, 211
 for DOA estimation, 297–298
 for frequency estimation, 174–175
 frequency selective, 194–203
 statistical accuracy of, 175
 estimate
 consistent, 91, 141, 154, 158, 159, 184, 185, 272, 291, 373
 statistically efficient, 375
 unbiased, 374
 Euclidean vector norm, 354
 exchange matrix, 362
 Expectation-Maximization algorithm, 188–194
 expected value, 5
 exponentially damped sinusoids, 253–256
 extended Rayleigh quotient, 350

F

far field, 276
 fast Fourier transform (FFT), 27–29
 for two-sided sequences, 21

 pruning in, 29
 radix-two, 27–29
 two-dimensional, 264, 269
 zero padding and, 29
 Fejer kernel, 32
 filter-bank methods, 217–232
 and periodogram, 220–222, 242–245
 APES, 256–259, 303
 for gapped data, 259–262
 for two-dimensional signals, 265–266
 refined, 222–232
 spatial APES, 319–326
 Fisher information matrix, 378
 flatness, spectral, 138
 forward prediction, 122, 123, 136
 forward-backward approach, 175–178
 frequency, 2, 3, 8
 angular, 3
 conversion, 3
 resolution, 33
 scaling, 16
 spatial, 284
 frequency band, 195
 frequency estimation, 153–178
 ESPRIT method, 174–175
 forward-backward approach, 175–178
 frequency-selective ESPRIT, 194–203
 FRES-ESPRIT, 194–203
 high-order Yule-Walker method, 162–166
 Min-Norm method, 171–173
 modified MUSIC method, 170
 MUSIC method, 166–169
 nonlinear least squares, 157–162
 Pisarenko method, 169
 spurious estimates, 170
 two-dimensional, 203–209
 frequency-selective method, 194–203
 Frobenius norm, 355, 365, 367

G

GAPES method, 259–262
 gapped data, 259–262
 Gaussian distribution
 circular, 380, 387

Gaussian random variable
 circular, 80
 Cramér–Rao bound for, 378–383
 moment property, 35
generalized Akaike information criterion, 411
generalized inverse, 365
Gohberg–Semencul formula, 128–130
grating lobes, 337

H

Hadamard matrix product, 358, 392
Hamming window, 45
Hankel matrix, 362
 block, 118
Hanning window, 45
Heisenberg uncertainty principle, 72
Hermitian matrix, 344, 348–351
Hermitian square root, 358
hypothesis testing, 183

I

idempotent, 294, 354
impulse response, 20, 72, 220, 224–226, 277
in-phase component, 280
inconsistent linear equations, 366–370
information matrix, 378
interior-point methods, 135
inverse covariances, 249

J

Jensen’s inequality, 192, 395

K

Kaiser window, 44, 45
kernel, 343
Kronecker delta, 4
Kronecker product, 265, 266
Kullback–Leibler information metric, 405–406

L

Lagrange multiplier, 308–309, 315
leading submatrix, 357

least squares, 20, 109, 171, 302, 303, 320, 329
 spectral approximation, 18
 with quadratic constraints, 308
least-squares method, 94, 96–99, 238, 255, 257, 259, 260, 263, 267, 268
least-squares solution, 367
Levinson–Durbin Algorithm, 100–101
 split, 102–106
likelihood function, 192, 375, 376, 378, 398
line spectrum, 153
linear equations
 consistent, 364–366
 inconsistent, 366–370
 least-squares solution, 367
 minimum–norm solution, 365
 systems of, 364–370
 total-least-squares solution, 369
linear prediction, 96, 122, 123, 125, 136–138
linear predictive modeling, 96
linearly parameterized basis, 203–209
low-pass signal, 278

M

MA covariance parameterization, 133
MA parameter estimation, 131–135
MA process, *see* moving average process
majorization, 190–191
majorizing function, 190
matrix
 centrosymmetric, 178, 332
 Cholesky factor, 358
 condition, 212, 366
 eigenvalue decomposition, 344–351
 exchange, 362
 fraction, 143
 Frobenius norm, 355, 365, 367
 Hankel, 362
 idempotent, 294, 354
 inversion lemma, 363
 Moore–Penrose pseudoinverse, 365
 orthogonal, 345
 partition, 359, 363
 persymmetric, 178, 332

- matrix (*continued*)
 - positive (semi)definite, 357–361
 - QR decomposition, 368
 - rank, 343
 - rank deficient, 343
 - semiunitary, 345, 349
 - singular value decomposition, 118, 164, 351–356
 - square root, 333, 358
 - Toeplitz, 362
 - trace, 346
 - unitary, 164, 174, 212, 345, 348, 351, 360, 368
 - Vandermonde, 361
- matrix fraction description, 143
- matrix inversion lemma, 258, 363
- maximum a posteriori detection, 402–405
- maximum likelihood estimate, 79, 157, 375, 382, 393, 398
 - of covariance matrix, 331–333
- maximum likelihood estimation, 398–402
 - regularity conditions, 400
- maximum likelihood method, 192
- MDL principle, 416
- mean-square convergence, 179
- mean squared error, 30
- Min–norm
 - and ESPRIT, 213
- Min–Norm method
 - and ESPRIT, 212
 - for DOA estimation, 297
 - for frequency estimation, 171–173
 - root, 172
 - spectral, 171
- minimization
 - cyclic, 189–190
 - majorization, 190–191
 - quadratic, 371–372
 - relaxation algorithms, 190
- minimum description length, 416
- minimum–norm constraint, 298–300
- minimum–norm solution, 365
- minimum phase, 93, 135
- missing data, 259–262
- model order selection, 375–376, 397–419
 - AIC rule, 409–412
 - BIC rule, 413–417
 - corrected AIC rule, 412
 - generalized AIC rule, 412–413
 - generalized information criterion, 411
 - Kullback–Leibler metric, 405–406
 - maximum a posteriori, 402–405
 - MDL rule, 416
 - multimodel, 418
- modified MUSIC method, 170, 203–209
- modified Yule–Walker method, 108–111
- modulation, 279
- Moore–Penrose pseudoinverse, 304, 365
- moving-average noise, 211
- moving-average parameter estimation, 131–135
- moving-average process
 - covariance structure, 93
 - definition of, 92
 - parameter estimation, 131–135
 - reflection coefficients of, 140
- moving-average spectral estimation, 106–107, 131–135
- multimodel order selection, 418
- multiple signal classification, *see* MUSIC method
- multivariate systems, 113–122
- MUSIC method
 - for DOA estimation, 297
 - modified, 170, 340
 - root, 169
 - spectral, 168
 - subspace-fitting interpretation, 339
- N
 - narrowband, 283
 - nilpotent matrix, 130
- NLS method, *see* nonlinear least squares method
- noise
 - complex white, 34
- noise gain, 312–314

- nonlinear least-squares method
 - for direction estimation, 294–295
 - for frequency estimation, 157–162
- nonsingular, 343
- normal equations, 95
- null space, 167, 343
- null-space basis, 203–209
- Nyquist sampling theorem
 - spatial, 285
 - temporal, 9

O

- order selection, *see* model order selection
- orthogonal complement, 353
- orthogonal matrix, 345
- orthogonal projection, 168, 198, 199, 354
- overdetermined linear equations, 109, 364, 366–370

P

- Padé approximation, 142
- parameter estimation
 - maximum likelihood, 398–402
- PARCOR coefficient, 101
- Parseval's theorem, 4, 132
- partial autocorrelation sequence, 122
- partial correlation coefficients, 101
- partitioned matrix, 359, 363
- periodogram
 - and frequency estimation, 159
 - bias analysis of, 30–34
 - definition of, 24
 - FFT computation of, 27–29
 - for two-dimensional signals, 263–264
 - properties of, 30–39
 - variance analysis of, 34–39
 - windowed, 50
- periodogram method, 24
- periodogram-based methods
 - Bartlett, 52–53
 - Daniell, 55–58
 - refined, 52–58
 - Welch, 53–55

- persymmetric matrix, 178, 332
- Pisarenko method
 - ARMA model derivation of, 210
 - for DOA estimation, 297
 - for frequency estimation, 166, 169
 - relation to MUSIC, 169
- planar wave, 276, 283
- positive (semi)definite matrices, 357–361
- postwindow method, 98
- power spectral density
 - and linear systems, 12
 - continuous, 90
 - definition of, 6, 7
 - properties of, 9
 - rational, 91
- prediction
 - backward, 122
 - forward, 136
 - linear, 96, 123, 125, 136–138, 386
- prediction error, 96, 386
- prewindow method, 98
- principal submatrix, 357
- probability density function, 72, 79, 375, 378
- projection matrix, 198, 199
- projection operator, 354

Q

- QR decomposition, 368
- quadratic minimization, 371–372
- quadratic program, 135
- quadrature component, 280

R

- random signals, 2
- range space, 167, 342
- rank, 199
- rank-deficient, 343
- rank of a matrix, 343
- rank of a matrix product, 343
- rational spectra, 91
- Rayleigh quotient, 349
 - extended, 350
- rectangular window, 45

reflection coefficient, 101
 properties of, 140
 region of convergence, 92
 RELAX algorithm, 190
 relaxation algorithms, 190
 resolution
 and time-bandwidth product, 72
 and window design, 42–44
 and zero padding, 29
 for filter-bank methods, 218
 for parametric methods, 162, 214
 frequency, 33
 limit, 33
 of beamforming method, 290, 334
 of Blackman–Tukey method, 41
 of Capon method, 236, 240, 249
 of common windows, 45
 of Daniell method, 57
 of periodogram, 23, 33
 of periodogram-based methods, 88
 spatial, 290, 334–337
 superresolution, 145, 146, 154
 Riccati equation, 116
 Rihaczek distribution, 16
 robust Capon method, 312–319
 root MUSIC
 for DOA estimation, 297
 for frequency estimation, 169
 row space, 342

S
 sample covariance, 24, 25, 53, 58, 75–77,
 80, 83
 sample covariance matrix
 ML estimates of, 331–333
 sampling
 Nyquist sampling theorem, 9
 spatial, 276, 285, 286
 temporal, 3
 semiparametric estimation, 326
 semidefinite quadratic program, 135
 semiunitary matrix, 345, 349
 sidelobe, 34, 44, 45
 signal modeling, 92

similarity transformation, 116, 175, 346
 singular value decomposition (SVD), 118,
 164, 304, 351–356
 sinusoidal signals
 amplitude estimation, 153
 ARMA model, 155
 covariance-matrix model, 156
 damped, 203–209, 253–256
 frequency estimation, 153–178
 models of, 150, 155–157
 nonlinear regression model, 155
 phase estimation, 153
 two-dimensional, 262–269
 skew-symmetric vector, 362
 Slepian sequences, 225–226
 two-dimensional, 265–266
 smoothed periodogram, 231
 smoothing filter, 137
 spatial filter, 287, 290
 spatial frequency, 284
 spatial sampling, 285, 286
 spatial spectral-estimation problem,
 275
 spectral analysis
 high-resolution, 154
 nonparametric, 2
 parametric, 2
 semi-parametric, 326
 superresolution, 154
 spectral density
 energy, 3
 power, 5
 spectral estimation
 definition of, 1, 13
 spectral factorization, 92, 132, 170
 spectral flatness, 138
 spectral-line analysis, 153
 spectral LS criterion, 131
 spectral MUSIC
 for DOA estimation, 297
 for frequency estimation, 168
 spectrum
 coherency, 13–15
 continuous, 90

- cross, 19
- discrete, 153
- rational, 91
- split Levinson algorithm, 102
- square root of a matrix, 333, 358
- stability
 - for AR models, 139
 - of AR estimates, 94
 - of Padé approximation, 142
 - of Yule–Walker estimates, 98, 139
- state-space equations
 - for ARMA process, 114
 - minimality, 117
 - nonuniqueness of, 116
- statistically efficient estimator, 375
- steering-vector uncertainty, 306–319
- structure indices, 114
- subarrays, 298
- submatrix
 - leading, 357
 - principal, 357
- subspace
 - and state-space representations, 114, 117–122
 - noise, 168
 - signal, 168
- superresolution, 145, 154
- symmetric matrix, 344
- symmetric vector, 362
- synthetic aperture, 334
- systems of linear equations, 364–370

T

- taper, 50
- Taylor series expansion, 373
- time width
 - definition of, 71
 - equivalent, 43, 53, 58, 73
- time-bandwidth product, 42–44, 70–75
- time-frequency distributions, 16
- Toeplitz matrix, 362
- total least squares, 109, 165, 171, 175, 369
- trace of a matrix, 346
- trace of a matrix product, 346

- transfer function, 11
- two-dimensional sinusoidal signals, 203–209
- two-dimensional spectral analysis
 - APES method, 267–269
 - Capon method, 267–269
 - periodogram, 263–264
 - refined filter-bank method, 265–266
- two-sided sequences, 21

U

- unbiased estimate, 374
- uncertainty principle, 71
- uniform linear array, 283–285
 - beamforming resolution, 334
 - spatial APES, 319–326
- unitary matrix, 164, 174, 212, 345, 348, 351, 360, 368

V

- Vandermonde matrix, 361
- vector
 - skew-symmetric, 362
 - symmetric, 362
- vectorization, 265, 268

W

- wave field, 275
 - planar, 276
- Welch method, 53–55
- white noise
 - complex, 34
 - real, 39
- whitening filter, 151
- Wiener–Hopf equation, 20
- window function
 - Bartlett, 45
 - Chebyshev, 44
 - common, 44–45
 - data and frequency-dependent, 62
 - design of, 42
 - Hamming, 45
 - Hanning, 45
 - Kaiser, 44
 - leakage, 34

window function (*continued*)

- main lobe, 33
- rectangular, 45
- resolution, 33
- resolution–variance tradeoffs, 42–44
- sidelobes, 33

Y

Yule–Walker equations, 95

Yule–Walker method

- for AR processes, 94

for DOA estimation, 295

for frequency estimation, 162–166

modified, 108–111

overdetermined, 109

stability property, 98

Z

zero padding, 28–29

zeroes

extraneous, 299, 300

in ARMA model, 92, 112